arXiv:1301.2598v1 [hep-th] 11 Jan 2013

A Sinister Universe: Chiral gravitons lurking, and Lyth un-bound

Peter Adshead,[1] Emil Martinec,[2] and Mark Wyman[1, 3]

[1]*Kavli Institute for Cosmological Physics, Enrico Fermi Institute,*
University of Chicago, Chicago, Illinois 60637, U.S.A
[2]*Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637, U.S.A.*
[3]*Department of Astronomy & Astrophysics, University of Chicago, Chicago, Illinois 60637, U.S.A.*

Models of inflation involving non-Abelian gauge field backgrounds can produce gravitational waves at an observable level with a preferred handedness, while satisfying current observational bounds. This asymmetry comes about because the non-Abelian background generates parity-violation in the action for perturbations. In the specific model we study, Chromo-Natural Inflation, these gravitational waves can be produced at observable levels even when no field makes a super-Planckian field excursion, thus evading a common formulation of the Lyth bound.

Scalar fields generically drive the inflationary epoch, however some of the qualitative problems of inflationary model building can be ameliorated by incorporating gauge fields into the dynamics [1–5]. In particular, non-Abelian gauge fields can appear in a homogeneous and isotropic configuration, leading to slow-roll inflation even on steep potentials [1] via competition between the potential forces and "magnetic drift" forces induced by a Chern-Simons interaction [5]. In this *Letter*, we study the perturbations in such models, and show that the gauge field background results in a parity-violating action for fluctuations. During inflation, this parity violation imprints on the spectrum of gravitational perturbations; one helicity is enhanced relative to the other, leading to a chiral spectrum of tensor fluctuations (chiral gravitational wave production was shown in a related context in [6]).

The amplitudes of the gravitational waves (in either polarization) is not directly correlated to the (sub-Planckian) excursion of the inflaton field during inflation, even though this is a "single-clock" inflationary setting, because the motion of the inflaton along its potential is mainly resisted by "magnetic drift," not Hubble friction. Thus, this mechanism evades the simplest version of the so-called "Lyth bound" [7]. A more comprehensive presentation of our analysis will appear in a forthcoming work [8].

Non-Abelian Gauge background: The Einstein-Yang-Mills equations are solved on FRW for a non-Abelian gauge field with this configuration:

$$A_0 = 0, A_i = \phi\delta^a_i J_a \equiv a\psi\delta^a_i J_a \tag{1}$$

$$F_{0i} = \partial_\tau\phi\delta^a{}_i J^a, F_{ij} = g\phi^2 f^a_{ij} J^a. \tag{2}$$

where J_a is a generator of SU(2) [9]. This gauge field background violates parity by relating space and gauge indices with a particular orientation. Perturbing around this background, one finds parity-violating interactions. We focus mainly on the particular case of Chromo-Natural Inflation (hereafter CNI), with $\mathcal{X}$ an axion, interacting with the gauge fields via

$$\mathcal{L} = \frac{1}{2}R - \frac{1}{2}(\partial\mathcal{X})^2 - V(\mathcal{X}) - \frac{1}{2}\mathrm{Tr}[F^2] - \frac{\lambda}{4f}\mathcal{X}\mathrm{Tr}[F\wedge F]. \tag{3}$$

In [1] we showed that this model inflates, with slow-roll provided via a magnetic-drift type force mediated by the Chern-Simons interaction. In the large drift force limit ($\lambda \gg 1$), the slow-roll equations for this model are

$$\dot{\psi} = -H\psi + \frac{f}{3g\lambda}\frac{V_{,\mathcal{X}}}{\psi^2}\ , \quad \frac{\lambda}{f}\dot{\mathcal{X}} = 2g\psi + \frac{2H^2}{g\psi}\ , \tag{4}$$

where g is the gauge coupling and f is the axion decay constant. Below we will take $V = \mu^4(1 + \cos(\mathcal{X}/f))$.

Perturbations: Before delving into the details, let us sketch the novel features of the perturbations in this scenario. Because we have a non-Abelian gauge field on a background, we will have additional propagating scalar, vector, and tensor modes. The perturbative dynamics exhibits standard behavior at early times $-k\tau \to \infty$, where one recovers free plane waves; also, for late times $k\tau \to 0$, the fluctuations freeze out, leaving only a single growing scalar and a single growing tensor mode. There are, however, two novel behaviors at intermediate times:

1. For $\boldsymbol{\lambda} m_\psi \gtrsim -(k\tau) \gtrsim m_\psi$, where $\boldsymbol{\lambda} = \lambda\psi/f$, and $m_\psi = g\psi/H$ is a characteristic mass scale of gauge fluctuations, we find a complex interacting system for the scalars. The mode that governs the amplitude of the axion at superhorizon scales decays by a factor of order $\boldsymbol{\lambda}$.

2. For a period $k\delta\tau \approx m_\psi$ around $-(k\tau) \approx 1$, one polarization of the gauge field tensor modes becomes unstable and grows exponentially. This mode sources the corresponding gravitational tensor polarization, enhancing it relative to the other polarization.

For the present work, we will ignore the vector perturbations, which do not play an important role.

Some Details: We will work with the ADM metric,

$$ds^2 = -N^2 d\tau^2 + a^2\tilde{h}_{ij}(dx^i + N^i dt)(dx^j + N^j dt) \tag{5}$$

where N is the lapse, N^i is the shift vector and $\tilde{h}_{ij}$ is the metric on spatial hypersurfaces. We choose spatially flat gauge, in which (with γ_{ij} transverse-traceless)

$$\tilde{h}_{ij} = a^2\left[e^{\gamma}\right]_{ij} = a^2\left[\delta_{ij} + \gamma_{ij} + \frac{1}{2}\gamma_{im}\gamma_{mj} + \dots\right]. \quad (6)$$

The gauge field perturbations decompose as

$$\delta A_\mu \equiv \Psi_\mu = (t^a_i + \epsilon^a_{ik}\chi_k + \delta^a_i\delta\phi)J_a\ , \quad (7)$$

(here t is symmetric and traceless) and the axion perturbs to $\mathcal{X} + \delta\mathcal{X}$. In what follows, we work with explicit components of the fields by choosing the wavenumber along the x^3 direction. The gauge modes (7) then have an SVT decomposition in which $t^{\pm} = \frac{1}{2}(t_{11} - t_{22}) \pm it_{12}$ form a transverse traceless tensor, $t_{3,(1\pm i2)}$ and $\chi_{1\pm i2}$ are transverse vectors, and $z \equiv \frac{1}{6}(2t_{33} - t_{11} - t_{22})$, χ_3 are scalars along with $\delta\phi$.

Tensors: The quadratic action for tensor modes is:

$$\begin{aligned}\mathcal{S}_T = \frac{1}{2}\int\frac{d^3k}{(2\pi)^3}d\tau\Big[&2\partial_\tau t^{\pm}_k\partial_\tau\bar{t}^{\pm}_k - 2\Big(k^2 + g\phi\frac{\lambda}{2f}\partial_\tau\mathcal{X}\Big)t^{\pm}_k\bar{t}^{\pm}_k\\ &\pm 2k\Big(\frac{\lambda}{f}\partial_\tau\mathcal{X} + 2g\phi\Big)t^{\pm}_k\bar{t}^{\pm}_k - 2\dot{\phi}(\partial_\tau t^{\pm}_k\bar{\gamma}^{\pm}_k + \partial_\tau\bar{t}^{\pm}_k\gamma^{\pm}_k)\\ &\mp 2kg\frac{\phi^2}{a}(t^{\mp}_k\bar{\gamma}^{\pm}_k + \bar{t}^{\pm}_k\gamma^{\pm}_k) + 2g^2\frac{\phi^3}{a}(t^{\pm}_k\bar{\gamma}^{\pm}_k + \bar{t}^{\pm}_k\gamma^{\pm}_k) \quad (8)\\ &+\frac{1}{2}\partial_\tau\gamma^{\pm}_k\partial_\tau\bar{\gamma}^{\pm}_k - \frac{1}{2}\Big(k^2 - \frac{1}{a}\frac{\partial^2 a}{\partial\tau^2} - 2\dot{\phi}^2 + 2g^2\frac{\phi^4}{a^2}\Big)\gamma^{\pm}_k\bar{\gamma}^{\pm}_k\Big],\end{aligned}$$

where again $\gamma_{\pm}$ are the transverse traceless gravitational modes. The canonically normalized fields are thus $\hat{\gamma}^{\pm} = \gamma^{\pm}/\sqrt{2}$ and $\hat{t}^{\pm} = \sqrt{2}t^{\pm}$. The crucial helicity dependence comes in the first term on the second line, which generates a tachyonic mass over a range of scales near horizon crossing for t^+ but not for t^-, and leads to a substantial growth of the former near horizon crossing. Following this growth, the coupling between the gravitational and gauge modes (which also has a mild chirality dependence) feeds the disparate amplitudes of the gauge tensor modes into the gravitational sector. It is worth emphasizing that this chiral splitting of tensor mode amplitudes is *not* solely due to the Chern-Simons interaction. The helicities are still split even in the absence of the Chern-Simons interaction ($\lambda \to 0$), because of the preferred orientation implicit in the gauge field background. Any non-Abelian model relying on a non-trivial gauge texture of the form (1) will produce a chiral spectrum of tensor modes [10].

Working in the quasi de Sitter limit where $H \approx$ const, $a \approx -1/H\tau$, and $\dot{\phi} \approx H\phi$, and introducing the variable $x = -k\tau$, the equation of motion for the gravitational wave can be written[11]

$$\begin{aligned}\hat{\gamma}^{\pm\prime\prime}_k + \Big(1 - \frac{2}{x^2} - 2\frac{\psi^2}{x^2} + 2\frac{g^2\psi^4}{H^2x^2}\Big)\hat{\gamma}^{\pm}_k& \quad (9)\\ =2\frac{\psi}{x}\hat{t}^{\pm\prime}_k \mp 2\frac{g\psi^2}{Hx}\hat{t}^{\pm}_k + 2\frac{g^2\psi^3}{H^2x^2}\hat{t}^{\pm}_k&\end{aligned}$$

where here and throughout primes denote derivatives w.r.t. x. The gauge field satisfies the approximate equation

$$\hat{t}^{\pm\prime\prime}_k + \left(1 + \frac{m}{x^2} \mp \frac{m_t}{x}\right)\hat{t}^{\pm}_k = 0. \quad (10)$$

The equation of motion for $\hat{t}^+$ has an exact solution

$$\hat{t}_k(x) = A_k M_{\alpha,\beta}(2ix) + B_k W_{\alpha,\beta}(2ix) \quad (11)$$

where $M_{\alpha,\beta}(2ix)$ and $W_{\alpha,\beta}(2ix)$ are the Whittaker M and W functions, and we have defined

$$m = m_\psi\frac{\lambda}{f}\frac{\dot{\mathcal{X}}}{H} = 2(1 + m^2_\psi) = \frac{1}{4} - \beta^2 \quad (12)$$

$$m_t = \frac{\lambda}{f}\frac{\dot{\mathcal{X}}}{H} + 2m_\psi = 2(1 + 2m^2_\psi)/m_\psi = -2i\alpha\ . \quad (13)$$

We set A_k and B_k, by demanding that the solutions approach canonically normalized positive frequency free plane waves as $x \to \infty$: $\hat{\gamma}^{\pm}, \hat{t}^{\pm} \to e^{ix}/\sqrt{2k}$. We find

$$A_k = \frac{(2i)^\alpha\mathcal{C}}{\Gamma(2\beta + 1)},\ B_k = \frac{2^\alpha i^{\beta+1}(-i)^{\alpha+\beta}\mathcal{C}}{\Gamma(\alpha + \beta + \frac{1}{2})} \quad (14)$$

where $\mathcal{C} = \Gamma(-\alpha + \beta + \frac{1}{2})/\sqrt{2k}$. We can now solve for the resulting positive helicity gravitational wave mode by inverting Eqn. (9). The resulting integrals can be evaluated in closed form in terms of Meijer G-functions. The late time result of this approximation scheme is

$$\hat{\gamma}^+(x) = \hat{\gamma}_0(x) + 2\psi B_k\Big(I_1 + m_\psi\left(I_2 - m_\psi I_3\right)\Big) \quad (15)$$

where $\hat{\gamma}_0(x)$ is the homogeneous solution and

$$\begin{aligned}I_1 =& -i\frac{\left(m^2 - 2imm_t + 2m - 2m_t^2\right)\sec(\pi\beta)\sin(\pi\alpha)\Gamma(\alpha)}{2m(m+2)}\\ &- i\frac{\pi^2\left(m^2 + 2imm_t + 2m - 2m_t^2\right)\sec(\pi\beta)\csc(\pi\alpha)}{2m(m+2)\Gamma(\alpha+1)\Gamma\left(\frac{1}{2} - \alpha - \beta\right)\Gamma\left(\frac{1}{2} - \alpha + \beta\right)},\\ I_2 =& \frac{\pi\sec(\pi\beta)\Gamma(-\alpha)}{2\Gamma\left(\frac{1}{2} - \alpha - \beta\right)\Gamma\left(\frac{1}{2} - \alpha + \beta\right)} + \frac{\pi(m - im_t)\sec(\pi\beta)}{2m\Gamma(1-\alpha)}\\ &- \frac{\pi\sec(\pi\beta)\Gamma(1-\alpha)}{m\Gamma\left(\frac{1}{2} - \alpha - \beta\right)\Gamma\left(\frac{1}{2} - \alpha + \beta\right)},\\ I_3 =& i\frac{\pi^2(m + im_t)\sec(\pi\beta)\csc(\pi\alpha)}{m(m+2)\Gamma(\alpha)\Gamma\left(\frac{1}{2} - \alpha - \beta\right)\Gamma\left(\frac{1}{2} - \alpha + \beta\right)}\\ &+ \frac{\pi(m_t + im)\sec(\pi\beta)}{m(m+2)\Gamma(-\alpha)}\ . \quad (16)\end{aligned}$$

We note that in the region where the approximation at (10) is breaking down (see the upper panel of Fig. 1), the metric fluctuations have frozen out and as a result, this approximation is excellent [12].

Scalar perturbations: We must also give a brief account of how scalar perturbations behave in this theory. A suitable gauge choice is non-Abelian Coulomb gauge

$$\bar{D}_i\Psi^i = \partial_i\Psi^i - ig[A_i, \Psi^i] = 0 \quad (17)$$



FIG. 1. The upper panel, dashed lines show the numerical integration of the equations resulting from the action, Eqn. (8). Solid lines show the result of the approximation of Eqns. (9) and (10). The values of the parameters are chosen so that $m_\psi \approx 2.1$ and $\psi \approx 0.048$. The lower panel shows the ratio of the power in the two gravitational wave helicities.

which results in a Gauss law constraint that is algebraic in k-space. This gauge condition can be used to eliminate χ_3 and, combined with the solution to the Gauss law constraint, leads to an action of the form

$$\mathcal{S}_S = \frac{1}{2}\int \mathcal{G}_{IJ}\Phi'_I\Phi'_J + \mathcal{M}_{IJ}\Phi_I\Phi_J + \mathcal{A}_{IJ}\Phi_I\Phi'_J \qquad (18)$$

in terms of fields $\Phi = (\hat{X}, \delta\phi, z)$, with $\hat{X} = a\,\delta\mathcal{X}$ the canonically normalized axion fluctuation; here the coefficient functions $\mathcal{G}$, $\mathcal{M}$ and $\mathcal{A}$ are rational functions of x with coefficients that depend on m_ψ and $\boldsymbol{\lambda}$. The action is that of a charged particle moving in three dimensions in a time-dependent harmonic well and magnetic field. The important terms in the action at large $\boldsymbol{\lambda}$ and x are a potential term $\boldsymbol{\lambda}^2 m_\psi^2 \hat{X}^2/x^2$ and "magnetic" coupling $(\boldsymbol{\lambda} m_\psi/x)(z-\delta\phi)\hat{X}'$. In the asymptotic past $x \to \infty$, the mass matrix $\mathcal{M} \to \mathbb{1}$, and $\mathcal{A} \to 0$, yielding free field equations for the modes. For $x \lesssim \boldsymbol{\lambda} m_\psi$, the mode of interest is a 'magnetic drift' mode where (as in the solution for the background) the large potential term balances against the large "magnetic" coupling, and the $\hat{X}$ kinetic term is largely irrelevant. A simplification of the dynamics ensues if we drop the $\hat{X}$ kinetic term altogether, as well as the subleading $O(\boldsymbol{\lambda}^0)$ terms in $\mathcal{M}_{\hat{x}\hat{x}}$. The axion is then determined in terms of the gauge scalars as

$$\hat{X} = \frac{(12m_\psi^4 + 6m_\psi^2 x^2 + x^4)\delta\phi + 2x^4 z - 3m_\psi^2 x(2m_\psi^2 + x^2)\delta\phi'}{\boldsymbol{\lambda} m_\psi^3 x^2}. \qquad (19)$$

Substituting this solution back into the action and making the further substitutions

$$\delta\phi \to \frac{x(2m_\psi\,\boldsymbol{\varphi} - x\,\boldsymbol{z})}{3(2m_\psi^2 + x^2)}\ ,\ z \to \frac{4m_\psi x\,\boldsymbol{\varphi} + (6m_\psi^2 + x^2)\boldsymbol{z}}{6(2m_\psi^2 + x^2)}\ , \qquad (20)$$

we find a simple reduced action depending only on m_ψ

$$\begin{aligned}\mathcal{S}_{red} = \int \Big[& \bar{\boldsymbol{\varphi}}'_k\boldsymbol{\varphi}'_k + \bar{\boldsymbol{z}}'_k\boldsymbol{z}'_k - \frac{2}{\sqrt{3}m_\psi}(\bar{\boldsymbol{z}}_k\boldsymbol{\varphi}'_k + \bar{\boldsymbol{\varphi}}_k\boldsymbol{z}'_k) \\ & - \frac{(m_\psi^2 - 2)}{{m_\psi}^2}\bar{\boldsymbol{\varphi}}_k\boldsymbol{\varphi}_k - \frac{2(m_\psi^2 + 1)}{\sqrt{3}m_\psi x}(\bar{\boldsymbol{z}}_k\boldsymbol{\varphi}_k + \bar{\boldsymbol{\varphi}}_k\boldsymbol{z}_k) \\ & - \frac{\left(6m_\psi^2 + x^2 + 6\right)}{3x^2}\bar{\boldsymbol{z}}_k\boldsymbol{z}_k\Big]\ .\end{aligned}$$

Note that the $\boldsymbol{\varphi}$ dynamics is unstable when $m_\psi^2 \lesssim 2$. In contrast to the tensor gauge modes, this scalar mode instability is present for a long time $m_\psi \lesssim x \lesssim \boldsymbol{\lambda} m_\psi$ and fully destabilizes the dynamics. There is thus a narrow range of m_ψ that could be consistent with data: small m_ψ leads to unstable scalar modes, while large m_ψ leads to unacceptably large tensor modes.

The amplitude of the axion on super-horizon scales is determined (up to order one corrections) by the behavior of $\boldsymbol{\varphi}$ near $x = 0$:

$$\hat{X} \sim \frac{\sqrt{2}\,\boldsymbol{\varphi}(0)}{\boldsymbol{\lambda}\,x} + O(x)\ . \qquad (21)$$

The limiting value $\boldsymbol{\varphi}(0)$ is of order one for large m_ψ, starts growing around $m_\psi \approx \sqrt{6}$, and diverges as $m_\psi \to \sqrt{2}$ due to the softening potential for this mode.

We can now estimate the amplitude of scalar and tensor fluctuations. In spatially flat gauge, the curvature perturbation is $\mathcal{R} = H\delta u$, where δu is the perturbation to the velocity potential, determined from

$$\begin{aligned} T_{0i} \approx\ & (\bar{p}_{\mathcal{X}} + \bar{p}_{\mathrm{YM}})a^2 N_i + a\dot{\mathcal{X}}\partial_i\delta\mathcal{X} \qquad (22) \\ & - \left(H\psi\partial_i(2z + 4\delta\phi) - g\psi^3\frac{\lambda}{f}\partial_i\delta\mathcal{X}\right) - 2g^2\frac{\phi^4}{a^2}N_i\ .\end{aligned}$$

The term $\propto g\psi^3\lambda/f$ is much larger than the rest. It comes from the gauge field momentum

$$T_{0i} = \mathrm{Tr}(E\times B)_i \approx g^2\mathrm{Tr}([A_j,\Psi_0][A_j,A_i]) = g^2\psi^3\Psi_0^i \qquad (23)$$

where Ψ_0^i is determined by the Gauss law constraint

$$\nabla\cdot E + \frac{\lambda}{f}\partial\mathcal{X}\cdot B = 0\ . \qquad (24)$$

At late times, the leading piece of the first term comes from $g^2[A_i,[A_i,\Psi_0]]$, and the leading piece of the second term is $(g\lambda/f)(\partial_i\delta\mathcal{X})[A_j,A_k]\epsilon^{ijk}$; thus one finds

$$\Psi_0^i = \frac{\lambda}{gf}\partial_i\delta\mathcal{X} \ , \tag{25}$$

leading to the relevant term in (22). The scalar curvature perturbation that results is

$$\mathcal{R} \approx \frac{H}{\bar\rho+\bar p} g\psi^3 \frac{\lambda}{f}\frac{\hat X}{a} = \frac{g\psi^3}{2H\epsilon_H}\frac{\lambda}{f}\frac{\hat X}{a} \ , \tag{26}$$

where $\epsilon_H = -\dot H/H^2$ and an overdot denotes a derivative with respect to cosmic time. The background has $\psi^3 \simeq V_{,\theta}/(3g\lambda H)$ and $\dot H \simeq V_{,\theta}\,\dot{\mathcal{X}}/(6fH)$, where $V_{,\theta}$ is the derivative of V w.r.t. $\theta = \mathcal{X}/f$, thus

$$\mathcal{R} \approx \frac{\delta\mathcal{X}}{\dot{\mathcal{X}}/H} \ . \tag{27}$$

Thus the axion is acting as the clock, as anticipated in [1], and the perturbation analysis is simply finding the magnitude of this perturbation. Now, relative to usual inflationary result for scalars on de Sitter, the magnitude of the $\delta\mathcal{X}$ perturbation is reduced from H by a factor of $\boldsymbol{\lambda}$; also, the axion velocity is reduced by a compensating factor due to "magnetic drift".

Putting it all together, one finds the scalar curvature fluctuation is given by

$$\mathcal{R} \simeq \frac{1}{\sqrt{2k^3}}\cdot\frac{Hm_\psi\boldsymbol{\varphi}(0)}{\sqrt{2}(1+m_\psi^2)}\Big(\frac{\lambda m_\psi V}{V_{,\theta}}\Big)^{1/2} \ ; \tag{28}$$

(note that in the reduced system, the scalar d.o.f.s are normalized to unit amplitude in the far past); expressed in terms of the slow roll parameters

$$\epsilon_H = -\frac{\dot H}{H^2} = -\frac{(1+m_\psi^2)V_{,\theta}}{\lambda m_\psi V} \tag{29}$$

$$\eta_H = \epsilon_H + \frac{1}{2}\frac{\dot\epsilon_H}{H\epsilon_H} = -\frac{(m_\psi^2+1)V_{,\theta\theta}}{\lambda m_\psi V_{,\theta}} - \frac{(m_\psi^2-1)V_{,\theta}}{\lambda m_\psi V} \ ,$$

one has

$$\mathcal{R} \simeq \frac{1}{\sqrt{2k^3}}\cdot\frac{H}{\sqrt{2\epsilon_H}}\cdot\frac{m_\psi\boldsymbol{\varphi}(0)}{(1+m_\psi^2)^{1/2}} \ . \tag{30}$$

Relative to the corresponding expressions for single-field inflation, the slow-roll parameters (29) directly exhibit the additional suppression of motion along the inflaton potential by the coupling λ to the gauge sector; and there is an additional enhancement from $\boldsymbol{\varphi}(0)$ when m_ψ is small enough. Differentiating (30) with respect to $dN = -Hdt$, one finds the spectral index

$$n_s - 1 \simeq -2\epsilon_H + \eta_H + d\log\boldsymbol{\varphi}(0)/dN \ . \tag{31}$$

The tensor-to-scalar ratio for the unenhanced graviton is

$$r_- \simeq \frac{8(1+m_\psi^2)\epsilon_H}{m_\psi^2\boldsymbol{\varphi}(0)^2} \ , \tag{32}$$

while the other tensor-to-scalar ratio r_+ is enhanced by the effect discussed above, determined from eq. (15). These formulae demonstrate that it is possible to satisfy current observational bounds with this model, while producing observable gravitational waves even in the absence of the chiral enhancement of one helicity mode. The chiral enhancement will be observable but not catastrophic for $m_\psi \lesssim 2$ (see Fig. 1).

Acknowledgements: We thank Wayne Hu for useful discussions and David Seery for early collaboration. This work was supported in part by DOE grant DE-FG02-90ER-40560 and by the Kavli Institute for Cosmological Physics at the University of Chicago through grants NSF PHY-0114422 and NSF PHY-0551142 and an endowment from the Kavli Foundation and its founder Fred Kavli.

Note Added: While we were writing up our results, parallel work appeared [13] independently deriving the stability condition $m_\psi^2 > 2$ using a WKB method.

---

[1] P. Adshead and M. Wyman, Phys.Rev.Lett., **108**, 261302 (2012), arXiv:1202.2366 [hep-th].

[2] P. Adshead and M. Wyman, Phys.Rev., **D86**, 043530 (2012), arXiv:1203.2264 [hep-th].

[3] A. Maleknejad and M. Sheikh-Jabbari, Phys.Rev., **D84**, 043515 (2011), arXiv:1102.1932 [hep-ph].

[4] M. M. Anber and L. Sorbo, Phys.Rev., **D81**, 043534 (2010), arXiv:0908.4089 [hep-th].

[5] E. Martinec, P. Adshead, and M. Wyman, (2012), arXiv:1206.2889 [hep-th].

[6] L. Sorbo, JCAP, **1106**, 003 (2011), arXiv:1101.1525 [astro-ph.CO].

[7] D. H. Lyth, Phys.Rev.Lett., **78**, 1861 (1997), arXiv:hep-ph/9606387 [hep-ph].

[8] P. Adshead, E. Martinec, and M. Wyman, *in prep.*

[9] Other embeddings of $SU(2) \subset SU(N)$ are possible; we take $N = 2$ for simplicity. We also take $\mathrm{Tr}[J_aJ_b] \equiv \frac{1}{2}\delta_{ab}$.

[10] The absence of parity violation in the effective action of ref. [14] indicates that some important effects have been missed. The premise of this work is that the gauge modes are heavy and may be integrated out for $m_\psi \gg 1$, while in fact there is in this limit a substantial period of time when certain gauge modes are not only light but unstable.

[11] To modify this analysis to encompass the gauge-flation model [3], simply replace $(\lambda/f)\partial_\tau\mathcal{X}/a$ with $\kappa H\epsilon_H g\psi^3/2$.

[12] Note, however, that the linearized equations for graviton production break down as m_ψ becomes large, as t^+ becomes exponentially large, and nonlinear effects such as $t^+t^+ \to \delta\mathcal{X}$ become important.

[13] E. Dimastrogiovanni and M. Peloso, (2012), arXiv:1212.5184 [astro-ph.CO].

[14] E. Dimastrogiovanni, M. Fasiello, and A. J. Tolley, (2012), arXiv:1211.1396 [hep-th].